

Lattices freely generated by posets within a variety.

Part I: Four easy varieties

Jean Yves Semegni and Marcel Wild

1 Introduction

This article constitutes the first of a two part essay triggered by [3]. Broadly speaking the intersection of Part I and Part II is the variety of distributive lattices. In Part II this variety is generalized to finitely generated lattice varieties, whereas in the present Part I three other "easy" varieties are taken aboard. These four varieties are

- the variety of all semilattices,
- the variety of all lattices,
- the variety of all distributive lattices,
- the variety of all Boolean lattices.

Strictly speaking the first and last are no lattice varieties, yet they fit well. These four varieties are easy in what concerns finding the free object generated by a poset. Some of the semilattice material will be useful in Part II. All results will be illustrated on a toy poset that accompanies us through both parts. Here comes the section break up.

An *implication* $A \rightarrow B$, where A, B are subsets of some fixed set P , is a certain Boolean formula with variables from P . In fact, it is the most frequently occurring kind of *Horn* formula. If Σ is a family of implications on P , then the family $\mathcal{C}(\Sigma)$ of all satisfying truth assignments, or more succinctly of all " Σ -closed" subsets $X \subseteq P$ is a closure system. Each closure system $\mathcal{C}$ is of type $\mathcal{C} = \mathcal{C}(\Sigma)$ for a suitable *implicational base* Σ . Section 2 outlines the (A, B) -algorithm [6] that computes $\mathcal{C}$ from Σ .

Unless stated otherwise, semilattice means $\vee$ -semilattice. According to Theorem 1 in section 3 each finite presentation $(P, \mathcal{R})$ of a semilattice S can be rewritten as a family Σ of implications on P , in which case S turns out to be isomorphic to $\mathcal{C}(\Sigma) \setminus \{\emptyset\}$. As a well known special case one obtains the semilattice S freely generated by a poset $(P, \leq)$, which is isomorphic to the semilattice of all nonempty order ideals of $(P, \leq)$ under union.

Section 4 and 5 are devoted to partial semilattices in general respectively particular. Partial semilattices $(P, \bigvee)$, that is, $\bigvee$ is a certain partial operation from P^ω to P , comprise posets $(P, \leq)$ as a special case, but are more specific than arbitrary presentations $(P, \mathcal{R})$ in that the free semilattice $S = F_{\bigvee}(P, \bigvee)$ generated by $(P, \bigvee)$ cannot collapse P . generalizing section 3, S can be viewed as a semilattice of certain *closed* order ideals of the underlying poset $(P, \leq)$.

As to the join core of a lattice L (section 5), according to Duquenne [1] it is the unique minimal partial semilattice $(P, \vee)$ such that $F_\vee(P, \vee) \simeq L$ as semilattices. We deemed it worthwhile to reprove this neat fact in a way that relates a bit more to the theory of implicational bases Σ .

Section 6 reviews, because it fits well, the lattice $FL(P, \leq)$ freely generated by a poset within the variety of all lattices.

The structure of the lattice $FD(P, \leq)$ freely generated by a poset within the variety $\mathcal{D}$ of all distributive lattices is also well known, but our proof in section 7 seems to be new.

Albeit as set, the variety $\mathcal{B}$ of all Boolean lattices is contained in $\mathcal{D}$, the lattice $FB(P, \leq)$ freely generated by $(P, \leq)$ within $\mathcal{B}$ is usually larger than $FD(P, \leq)$ because $FB(P, \leq)$ additionally needs to be closed under complementation. The structure of $FB(P, \leq)$ is determined by the number t of atoms. Finding t in terms of $(P, \leq)$ is easy but believed to be new. For our toy poset $(P, \leq)$ the cardinalities of $FL(P, \leq)$, $FD(P, \leq)$, and $FB(P, \leq)$ are shown to be 35, 25, and 16384 respectively.

2 Implications and the (A, B) -algorithm

An *implication* on a set P is a pair of subsets (A, B) , written as $A \rightarrow B$. Here A and B are the *premise* and *conclusion* of the implication. Let

$$\Sigma := \{A_1 \rightarrow B_1, \dots, A_n \rightarrow B_n\} \quad (1)$$

be a family of implications. A subset $X \subseteq P$ is Σ -closed if

$$(\forall 1 \leq i \leq n) (A_i \subseteq X \Rightarrow B_i \subseteq X) \quad (2)$$

In other words, for each i one must have either $A_i \not\subseteq X$ or $B_i \subseteq X$ (or both). It is easy to see that the family $\mathcal{C}(\Sigma)$ of all Σ -closed subsets is a *closure system*, that is, it contains P and is closed under intersections. Observe that the empty set $\emptyset$ is in $\mathcal{C}(\Sigma)$ if and only if there are no implications of type $\emptyset \rightarrow B_i$ ($B_i \neq \emptyset$). There is no need to deepen the mentioned connection (introduction) to Horn formulae here; that is done in [6].

Example 1 Let $P = \{a, b, c, d, e, f, g\}$ and

$$\Sigma := \{ \{b\} \rightarrow \{a, e\}, \{c\} \rightarrow \{b\}, \{d\} \rightarrow \{c, f\}, \\ \{f\} \rightarrow \{e\}, \{g\} \rightarrow \{b, f\}, \{b, f\} \rightarrow \{g\} \}$$

Then $X = \{a, b, e\}$ is Σ -closed since with $\{b\} \subseteq X$ also $\{a, e\} \subseteq X$. Ditto $\{a, e\}$ is Σ -closed, but not $\{a, b\}$. Here is the complete closure system:




Figure 1:

Let us give a rough sketch of the (A, B) -algorithm [6] which efficiently generates closure systems of the kind $\mathcal{C}(\Sigma)$.

Example 2

Let Σ be as in Example 1. The family r of all sets $X \in 2^P$ that satisfy $\{b\} \rightarrow \{a, e\}$ can compactly be written¹ as

	a	b	c	d	e	f	g
$r =$	β	α	2	2	β	2	2

That is, each 0, 1-incidence vector corresponding to such a set X must be such that if its second component α is 1, then also 1 must occur at the two positions labelled β . The other positions carry a label 2 which indicates that they are free to be independently 0 or 1. In order to impose the second implication $\{c\} \rightarrow \{b\}$ we first split r into the disjoint union of $r_1 = \{X \in r \mid b \notin X\}$ and $r_2 = \{X \in r \mid b \in X\}$. Thus

	a	b	c	d	e	f	g
$r_1 =$	β	α	0	2	β	2	2
$r_2 =$	β	α	1	2	β	2	2

All $X \in r_1$ satisfy $\{c\} \rightarrow \{b\}$ since $c \notin X$. But $X \in r_2$ satisfies $\{c\} \rightarrow \{b\}$ if and only if $\alpha = 1$ in r_2 . The latter entails that both β in r_2 are 1. Hence the family of all $X \in 2^P$ that satisfy both of $\{b\} \rightarrow \{a, e\}$ and $\{c\} \rightarrow \{b\}$ is

a	b	c	d	e	f	g
β	α	0	2	β	2	2
1	1	1	2	1	2	2

¹We used α, β instead of a, b in [6] because a and b have another meaning in the present article.

Continuing by imposing $\{d\} \rightarrow \{c, f\}$ up to $\{b, f\} \rightarrow \{g\}$ one obtains the table

a	b	c	d	e	f	g
β	α	0	0	β	0	0
2	0	0	0	1	1	0
1	1	0	0	1	1	1
1	1	1	0	1	0	0
1	1	1	2	1	1	1

which encodes $\mathcal{C}(\Sigma)$. For instance, the first row comprises the five sets $\phi, \{a\}, \{e\}, \{a, e\}$ and $\{a, b, e\}$. Since the rows comprise mutually disjoint set families, one obtains

$$|\mathcal{C}(\Sigma)| = 5 + 2 + 1 + 1 + 2 = 11$$

in accordance with Figure 1.

One can show that the wasteful deletion of rows during the (A, B) -algorithm can be avoided. Another nice feature is that the time spent is mainly dependent on the number of implications and *not* on the actual size of $\mathcal{C}(\Sigma)$. The (A, B) -algorithm simplifies to the (a, B) -algorithm, and runs particularly well, when all premises are singletons.

Back from algorithms to theory. As is well known, every closure system $\mathcal{C}$ on P (not necessarily induced by Σ) comes along with a closure operator $A \mapsto \overline{A}$ ($A \subseteq P$) defined by

$$\overline{A} := \bigcap \{X \in \mathcal{C} \mid X \supseteq A\}. \quad (3)$$

Furthermore $\mathcal{C}$, partially ordered by inclusion, is a $\vee$ -semilattice with joins (suprema) given by

$$A \vee B = \overline{A \cup B}. \quad (4)$$

Conversely, every closure operator $A \mapsto \overline{A}$ yields the closure system $\mathcal{C}$ of all *closed* sets $A = \overline{A}$. Given any closure system $\mathcal{C}$ on P , a family Σ of implications with $\mathcal{C}(\Sigma) = \mathcal{C}$ is called an *implicational base* of $\mathcal{C}$. It is *nonredundant* if no proper subset of Σ is an implicational base of $\mathcal{C}$. Any two nonredundant implicational bases Σ_1, Σ_2 of $\mathcal{C}$ yield the same set $E(\mathcal{C})$ of *essential* elements in that

$$\{\overline{A} \mid (A \rightarrow B) \in \Sigma_1\} = \{\overline{C} \mid (C \rightarrow D) \in \Sigma_2\} := E(\mathcal{C}). \quad (5)$$

An implicational base Σ is *optimal* if the sum of the cardinalities of all premises and conclusions of implications occurring in Σ is minimal. One can show that each optimal implicational base is nonredundant. A nonredundant implicational base can be computed in quadratic time, but computing an optimal implicational base is *NP*-hard. Interestingly, if $\mathcal{C}$ is modular as a lattice, the task can be achieved in polynomial time.

Along with each closure operator $X \mapsto \overline{X}$ on P comes the *quasi-closure* operator $X \mapsto X^\bullet$ which is defined by

$$X^\bullet := X \cup X^\circ \cup X^{\circ\circ} \cup X^{\circ\circ\circ} \cup \dots, \quad (6)$$

where

$$Y^\circ := Y \cup \bigcup \{ \overline{Z} \mid Z \subseteq Y \text{ and } \overline{Z} \neq \overline{Y} \}. \quad (7)$$

Notice that the iterated sets $X^{\circ \circ \dots \circ}$ in (6) eventually become stationary due to the finiteness of P . It is clear that $X \mapsto X^\bullet$ is a closure operator and that $X^\bullet \subseteq \overline{X}$ for all $X \subseteq P$. Call $X \subseteq P$ *quasiclosed* if $X^\bullet = X$.

3 Finitely presented semilattices

Let P be any finite set of “symbols” and let $\mathcal{R}$ be a finite set of semilattice relations with symbols from P . The *semilattice* $F_\vee(P, \mathcal{R})$ *freely generated by the set P and subject to the relations in $\mathcal{R}$* is the up to isomorphism unique² semilattice S such that

- (a) There is a map $\phi : P \rightarrow S$ which *satisfies* the relations from $\mathcal{R}$ (obvious definition) and is such that $\phi(P)$ generates S .
- (b) For each semilattice T and each map $\phi : P \rightarrow T$ *respecting* the relations from $\mathcal{R}$ (obvious definition), there is a $\vee$ -homomorphism $\Phi : F_\vee(P, \mathcal{R}) \rightarrow T$ that extends ϕ .

Each semilattice relation in $\mathcal{R}$, say $a \vee b = c \vee d$, can be rewritten as $a \vee b \geq c \vee d$ and $c \vee d \geq a \vee b$. Of course $\mathcal{R}$ can also feature “lone” inequalities $x \geq y$, since this amounts to $x \vee y = x$. So we may suppose that all relations in $\mathcal{R}$ are of type

$$a \vee b \vee \dots \vee c \geq d \vee e \vee \dots \vee f. \quad (8)$$

Let $\Sigma(\mathcal{R})$ be the family of all corresponding implications

$$\{a, b, \dots, c\} \rightarrow \{d, e, \dots, f\}. \quad (9)$$

Theorem 1 [5, Satz 45] *If all relations in $\mathcal{R}$ have been adjusted to type (8), then the semilattice $F_\vee(P, \mathcal{R})$ is isomorphic to the semilattice $\mathcal{C}(\Sigma(\mathcal{R})) \setminus \{\emptyset\}$, with joins as in (4).*

Proof: We first show that $S := \mathcal{C}(\Sigma(\mathcal{R})) \setminus \{\emptyset\}$ satisfies (a). For all $a \in P$ put $\overline{a} := \overline{\{a\}}$ and define $\phi(a) := \overline{a}$. That S is generated by $\overline{P} := \phi(P)$ can be seen from

$$\overline{\{a, b, \dots, c\}} = \overline{\overline{a} \cup \overline{b} \cup \dots \cup \overline{c}} \stackrel{(4)}{=} \overline{a} \vee \overline{b} \vee \dots \vee \overline{c}.$$

²Recall that unless stated otherwise, “semilattice” means $\vee$ -semilattice. The semilattice $F_\vee(P, \mathcal{R})$ is a special case of a universal algebra *finitely presented by generators and relations*. Such creatures are always unique up to isomorphism.

Let $a \vee b \vee \dots \vee c \geq d \vee e \vee \dots \vee f$ be a relation from $\mathcal{R}$. Since $\{a, \dots, c\} \rightarrow \{d, \dots, f\}$ belongs to $\Sigma(\mathcal{R})$, and $\overline{\{a, \dots, c\}}$ is $\Sigma(\mathcal{R})$ -closed, we conclude $\overline{\{a, \dots, c\}} \supseteq \{d, \dots, f\}$, which yields

$$\bar{a} \vee \dots \vee \bar{c} = \overline{\{a, \dots, c\}} \supseteq \overline{\{d, \dots, f\}} = \bar{d} \vee \dots \vee \bar{f}$$

This shows (a). As to (b), let T be any semilattice and let $\phi : P \rightarrow T$ respect $\mathcal{R}$. We claim:

$$\text{If } g \in \overline{\{a, b, \dots, c\}}, \text{ then } \phi(g) \leq \phi(a) \vee \phi(b) \vee \dots \vee \phi(c) \quad (10)$$

To fix ideas, say $\{a, b\} \rightarrow \{d\}$ and $\{c, d\} \rightarrow \{g\}$ belong to $\Sigma(\mathcal{R})$, and so $g \in \overline{\{a, b, c\}}$. Since ϕ respects the relation $a \vee b \geq d$, one has $\phi(a) \vee \phi(b) \geq \phi(d)$ in T . Ditto $\phi(c) \vee \phi(d) \geq \phi(g)$. But this yields $\phi(a) \vee \phi(b) \vee \phi(c) \geq \phi(g)$, which proves (10). Define $\Phi : S \rightarrow T$ by

$$\Phi(\overline{\{a, b, \dots, c\}}) := \phi(a) \vee \phi(b) \vee \dots \vee \phi(c)$$

Because of (10), Φ is well defined, i.e. not dependent on the particular generators $a, b, \dots, c$. From $\overline{\{a, \dots, c\}} \vee \overline{\{d, \dots, e\}} = \overline{\{a, \dots, c, d, \dots, e\}}$ readily follows that Φ is a $\vee$ -morphism. ■

We mention that Theorem 1 generalizes in natural ways to finitely presented commutative semigroups.

Example 3 By Theorem 1 and Example 1, if $P = \{a, b, \dots, g\}$, and $\mathcal{R}$ consists of the relations $b \geq a \vee e$, $c \geq b$, $d \geq c \vee f$, $f \geq e$, $g = b \vee f$, then




Figure 2:

Relations not present in $\mathcal{R}$ may well be present in $F_{\vee}(P, \mathcal{R})$, such as $e \leq c$. But such relations are always *deductible* from the relations in $\mathcal{R}$. Thus $e \leq c$ follows from $e \leq a \vee e \leq b \leq c$, where the first inequality follows from the semilattice axioms, and the other two are relations in $\mathcal{R}$. The relations in $\mathcal{R}$ may even cause the collapse of $F_{\vee}(P, \mathcal{R})$ to a single point, say if $\mathcal{R} = \{a \geq b, b \geq c, c \geq a\}$.

Let $(P, \leq)$ be a finite poset. The semilattice freely generated by $(P, \leq)$ is defined as

$$F_{\vee}(P, \leq) = F_{\vee}(P, \mathcal{R})$$

where $\mathcal{R}$ consists of all relations $a \geq b$ holding in P . Because $\vee$ does not feature in $\mathcal{R}$, the only possible deductions use the transitivity of $\geq$. But if $a \geq b$ and $b \geq c$ are in $\mathcal{R}$ then so is $a \geq c$. Hence $\mathcal{R}$ is deductively closed.

Corollary 1 *The semilattice $F_\vee(P, \leq)$ is isomorphic to the $\cup$ -semilattice $Id(P, \leq) \setminus \{\emptyset\}$. Furthermore, $F_\vee(P, \leq)$ is up to isomorphism the unique semilattice S such that:*

- (i) *There is a generating subset of S which as a poset is isomorphic to $(P, \leq)$.*
- (ii) *For each semilattice T and each monotone map $\phi : P \rightarrow T$ there is a semilattice morphism $\Phi : S \rightarrow T$ that extends ϕ .*

Proof: By Theorem 1, $F_\vee(P, \leq)$ is isomorphic to $\mathcal{C}(\Sigma(\mathcal{R})) \setminus \{\emptyset\}$ which of course is $Id(P, \leq) \setminus \{\emptyset\}$. Properties (i) and (ii) essentially follow from (a), (b) (beginning of sec.3). The fact that in (i) the generating set of S cannot collapse, but is *isomorphic* to P , is due to the fact that here $\mathcal{R}$ is deductively closed. ■

Mutatis mutandis, all holds when $\vee$ is switched with $\wedge$. Specifically, if $(P^d, \leq^d)$ is the dual of $(P, \leq)$, then

$$F_\wedge(P, \leq) \cong F_\vee(P^d, \leq^d) \cong Id(P^d, \leq^d) \setminus \{\emptyset\} \cong Fil(P, \leq) \setminus \{\emptyset\},$$

where $Fil(P, \leq)$ is the family of order filters of $(P, \leq)$.

Example 4 Along with the poset $(P, \leq)$ in Fig.3(a) the freely generated $\vee$ -semilattice $F_\vee(P, \leq)$ (Fig.3(b)) and the freely generated $\wedge$ -semilattice $F_\wedge(P, \leq)$ (Fig.3(c)) are shown.

Notice that despite first appearances $F_\wedge(P, \leq)$ is *not* the dual of $F_\vee(P, \leq)$. The above isomorphism $F_\wedge(P, \leq) \rightarrow Fil(P, \leq) \setminus \{\emptyset\}$ sends (say) c to $c\uparrow$ and f to $f\uparrow$, but $c \wedge f$ maps to $c\uparrow \cup f\uparrow$. Thus the semilattice operation in $Fil(P, \leq) \setminus \{\emptyset\}$ is $\cup$.

4 Partial semilattices

Every subset P of a semilattice S determines a partial operation $\bigvee$ from P^ω to P by setting $\bigvee X := \sup(X)$ if $\sup(X) \in P$. If $\sup(X) \in S \setminus P$, then $\bigvee X$ is not defined. One calls the pair $(P, \bigvee)$ a *partial semilattice*³. Let $\mathcal{R}$ be the set of semilattice relations that bijectively correspond to those subsets $X \subseteq P$ for which $\bigvee X$ is defined. Then $\mathcal{R}$ is deductively closed since any relation not in $\mathcal{R}$ but deducible from $\mathcal{R}$, would in particular

³An *intrinsic* definition of $(P, \bigvee)$, i.e. one that avoids a comprising semilattice S is possible. Essentially, each poset $(P, \leq)$ can be enriched to a partial semilattice by focussing on any subsets $X_i \subseteq P$ for which $\sup(X_i)$ happens to exist in $(P, \leq)$, and by defining the partial operation just for these X_i . That may force the definition of additional $\bigvee Y$ but will not lead to contradictions. Notice that [2] only considers *binary* partial operations. That would be too restrictive for us in the sequel.




Figure 3:

hold in S , but it doesn't. Gathering all relations of type $a \geq b$ in $\mathcal{R}$ yields a unique poset $(P, \leq)$. An order ideal A of $(P, \leq)$ is called $\vee$ -ideal if

$$(\forall X \subseteq A) \quad \left(\bigvee X \in P \Rightarrow \bigvee X \in A \right)$$

(Here of course " $\bigvee X \in P$ " is shorthand for "the partial operation $\bigvee$ is defined for X ".) One readily verifies that the set $Id(P, \bigvee)$ of all $\vee$ -ideals is a closure system.

Let $(P, \bigvee)$ be a partial semilattice. The semilattice *freely generated* by $(P, \bigvee)$ is defined as $F_v(P, \bigvee) := F_v(P, \mathcal{R})$ where $\mathcal{R}$ is as above.

Corollary 2 *Given a partial semilattice $(P, \bigvee)$ the semilattice $F_v(P, \bigvee)$ is isomorphic to $Id(P, \bigvee) \setminus \{\emptyset\}$ with joins given as in (4). Moreover, $F_v(P, \bigvee)$ is up to isomorphism the unique semilattice S such that*

- (A) *There is a generating subset of S which as a partial semilattice is isomorphic to $(P, \bigvee)$*
- (B) *For each semilattice T each partial morphism $\psi : P \rightarrow T$ (i.e. $\psi(\bigvee X) = \bigvee \psi(X)$ whenever $\bigvee X$ exists) can be extended to a semilattice morphism $\Phi : S \rightarrow T$.*

Proof: By Theorem 1, $F_v(P, \bigvee)$ is isomorphic to $\mathcal{C}(\Sigma(\mathcal{R})) \setminus \{\emptyset\}$ which clearly is $Id(P, \bigvee) \setminus \{\emptyset\}$. Properties (A) and (B) follow from (i), (ii) by taking into account that $\mathcal{R}$ is deductively closed (cf proof of Corollary 1). ■

5 The core of a semilattice

Each semilattice becomes a lattice by attaching a smallest element 0. In this article we exploit the converse, namely that each lattice L is in particular a semilattice, and that finite presentations of semilattices are easier to handle. All of the sequel owes a lot to [1]. More details at the end of this section.

For starters, for any lattice L , the semilattice $L \setminus \{0\}$ is isomorphic to a finitely presented semilattice $F_\vee(P, \mathcal{R})$ when P is taken as the set $J = J(L)$ of nonzero join irreducibles. As to $\mathcal{R}$, for $a \in L$ let $J(a) := \{p \in J \mid p \leq a\}$, and for $X \subseteq J$ define its "natural closure" as $\overline{X} := J(\bigvee X)$. If $\mathcal{C}$ is the associated closure system, then $a \mapsto J(a)$ yields an isomorphism from L to $\mathcal{C}$. If Σ is any implicational base of $\mathcal{C}$ and $\mathcal{R}$ the corresponding set of semilattice relations, then $L \setminus \{0\} \cong F_\vee(J, \mathcal{R})$ by Theorem 1.

Trouble is that Σ is not readily found, and this requires some detours. First, the set J becomes a poset $(J, \leq)$ with the ordering induced by L . With respect to the above closure system each closed set $X = J(a)$ is an order ideal, but (unless L is distributive) not every order ideal X of $(J, \leq)$ is closed. The trick will be to extend J to a suitable partial semilattice $(P, \vee)$ and to compute L as the closure system $Id(P, \vee)$ of all $\vee$ -ideals of $(P, \vee)$ (see Corollary 2).

Thus, given a lattice L , consider any subset $P \subseteq L$ for which

$$\psi : L \rightarrow Id\left(P, \bigvee\right), \quad a \mapsto P(a) := \{p \in P \mid p \leq a\} \quad (11)$$

is a semilattice isomorphism. For instance, $P := L$ does the job, but we strive for P to be as small as possible. One always needs $J = J(L) \subseteq P$. Indeed, if we had $q \notin P$ for some $q \in J$, then $\psi(q) = \psi(q^*)$ where q^* is the unique lower cover of q . This contradicts the injectivity of ψ . Conversely, for any superset P of J , the map ψ defined in (11) will be injective because $a \neq b$ implies $J(a) \neq J(b)$ which, in view of $J \subseteq P$, amounts to $P(a) \cap J \neq P(b) \cap J$, and hence $P(a) \neq P(b)$.

Assume again that ψ in (11) is an isomorphism. In order to derive a stronger necessary condition than just $J \subseteq P$, let us return to the quasi-closure of (6). For $q \in J$ one has $\{q\}^\bullet = \{q\}$ and hence $\{q\}^\bullet \neq \overline{\{q\}} = J(q)$ unless q is an atom of L . In particular, each *non-atomic* $q \in J$ is such that $J(q)$ contains a nonclosed quasiclosed generating set.

Consider now a *reducible* $a \in L$ such that $J(a)$ contains a quasiclosed nonclosed generating set K , that is, $K = K^\bullet \neq \overline{K} = J(a)$. We want to show that $a \in P$, and so by contraposition assume that $a \notin P$. By quasiclosedness K must be an order ideal of $(J, \leq)$. In order to see that

$$K_P := \left\{ \bigvee X \mid X \subseteq K, \bigvee X \in P \right\}$$

is an order ideal of $(P, \leq)$, fix $b \in P$ and $\bigvee X \in K_P$ with $b \leq \bigvee X$. From $\bigvee X < \bigvee K$ (since $\bigvee K = a \notin P$) follows $J(\bigvee X) \subseteq K$ since $K = K^\bullet$. Hence there is $Y \subseteq J(\bigvee X) \subseteq K$ with $\bigvee Y = b$, and so $b \in K_P$. In fact, K_P even is a $\bigvee$ -ideal of the partial semilattice $(P, \bigvee)$ because if $\bigvee X, \bigvee Y \in K_P$ with $c := (\bigvee X) \vee (\bigvee Y) \in P$, then $c = \bigvee (X \cup Y) \in K_P$ by definition of K_P . Hence K_P is a member of $Id(P, \bigvee)$. But we claim it is not in the range of ψ . Since $\bigvee K = a$, the only possibility for that to happen is $K_P = \psi(a)$. However, by assumption there is some $r \in J(a) \setminus K$, and so $r \in \psi(a) \setminus K_P$. This contradiction shows that for ψ to be an isomorphism, it is necessary that P comprises the *join core*

$$K_\vee(L) := J(L) \cup E_\vee(L), \quad (12)$$

where the set of *join-essential* elements is

$$E_\vee(L) := \{a \in L \mid (\exists K \subseteq J(a)) \ K^\bullet \neq \overline{K} = J(a)\} \quad (13)$$

Theorem 2 *If L is a finite lattice and $P := K_\vee(L)$, then $L \cong F_\vee(P, \bigvee)$ as semilattices. Furthermore, for each subset Q of L , one has $L \cong F_\vee(Q, \bigvee)$ if and only if $P \subseteq Q$.*

Proof: By the deliberations above it suffices to prove the first claim. Recall from Corollary 2 that $F_\vee(P, \bigvee) \simeq Id(P, \bigvee) \setminus \{\emptyset\}$. Because of $J \subseteq P$ we know that ψ in (11) is injective. In order to see that ψ is surjective (and thus clearly an isomorphism), we show that for each nonempty H in $Id(P, \bigvee)$ one has

$$H = P(a), \quad \text{where } a := \bigvee H \quad \left(= \bigvee (J \cap H) \right). \quad (14)$$

By contraposition, assume that H is an inclusion-minimal counter example of (14). If we had $a := \bigvee H \in P$, then $a \in H$ because H is a $\bigvee$ -ideal, and so $P(a) \subseteq H$, i.e. $P(a) = H$. Because by assumption $H \subsetneq P(a)$, we conclude that $a \notin P$. Thus we get a desired contradiction to (12), (13) if we can establish $J \cap H$ as a quasiclosed nonclosed subset of $J(a)$.

As to "quasiclosed", we need to show that

$$\left(X \subseteq J \cap H \text{ and } \bigvee X < a \right) \text{ implies } J\left(\bigvee X\right) \subseteq H.$$

Case 1: $\bigvee X \in P$. Then $\bigvee X \in H$, and so $J(\bigvee X) \subseteq P(\bigvee X) \subseteq H$.

Case 2: $\bigvee X \notin P$. Let K_P be the $\bigvee$ -ideal generated by X (recall that $Id(P, \bigvee)$ is a closure system). Thus $X \subseteq K_P \subseteq H$. In fact $K_P \subsetneq H$ since $\bigvee X < a$, $\bigvee H = a$. But then (14) holds for K_P by the minimality of H , and so $J(\bigvee X) \subseteq P(\bigvee X) = K \subseteq H$.

As to $J \cap H$ being nonclosed, let $b \in P(a) \setminus H$ be minimal. Suppose b was reducible. Then $b = \bigvee Q$ with $Q \subseteq J(b)$, and $J(b) \subseteq H$ by the minimality of b . This is impossible since $Q \subseteq H$, $\bigvee Q \in P$ would force $\bigvee Q \in H$. Therefore $b \in J(a) \setminus (J \cap H)$. ■

Duquenne [1] studies the *meet* core $K_\wedge(L)$ for various kinds of lattices. Up to duality his definition of $K_\wedge(L)$ matches our definition of the join core $K_\vee(L)$ in (12). However, Duquenne's $\wedge$ -essential elements by definition are the *reducible* members of $K_\wedge(L)$,

whereas we defined $E_\vee(L)$ in such a way that $E_\vee(L) \cap J(L) = \{q \in J(L) \mid q \text{ is no atom}\}$. In so doing $E_\vee(L)$ matches the natural definition of $E(\mathcal{C})$ in (a),(b) when $\mathcal{C} := \{J(a) \mid a \in L\}$. For instance, for a finite Boolean lattice L equation (12) becomes $K_\vee(L) = J(L) \cup \emptyset$, corresponding to the fact that only the *empty* family Σ is a *nonredundant* implicational basis of $\mathcal{C} := \{J(a) \mid a \in L\} \simeq 2^{J(L)}$.

6 The variety of all lattices

In this and the next two sections we turn from freely generated semilattices to freely generated lattices. Let t_1, t_2, t_3 be elements of any lattice. Then, clearly:

$$(i) \quad t_1 \vee t_2 \leq t_3 \quad \Leftrightarrow \quad t_1 \leq t_3 \quad \text{and} \quad t_2 \leq t_3$$

$$(ii) \quad t_3 \leq t_1 \wedge t_2 \quad \Leftrightarrow \quad t_3 \leq t_1 \quad \text{and} \quad t_3 \leq t_2$$

$$(iii) \quad t_3 \leq t_1 \quad \text{or} \quad t_3 \leq t_2 \quad \Rightarrow \quad t_3 \leq t_1 \vee t_2$$

$$(iv) \quad t_1 \leq t_3 \quad \text{or} \quad t_2 \leq t_3 \quad \Rightarrow \quad t_1 \wedge t_2 \leq t_3$$

Definition: The *lattice freely generated by the poset P* within a variety $\mathcal{V}$ of lattices is defined as the up to isomorphism unique lattice $F\mathcal{V}(P, \leq)$ in $\mathcal{V}$ which satisfies:

- (a) $F\mathcal{V}(P, \leq)$ is generated by P
- (b) For each lattice $T \in \mathcal{V}$ and each monotone map $\phi : P \rightarrow T$, there is a homomorphism $\Phi : F\mathcal{V}(P, \leq) \rightarrow T$ that extends ϕ .

In this section $\mathcal{V}$ is the variety of *all* lattices and we write $FL(P, \leq)$ for $F\mathcal{V}(P, \leq)$. It turns out⁴ that the converse implication in (iii) respectively (iv) holds in every lattice $FL(P, \leq)$.

For instance, P being the poset from figure 3(a), is it true that $b \vee (c \wedge f) \leq c \wedge (a \vee f)$ in $FL(P, \leq)$? By (i) the truth amounts to

$$b \leq c \wedge (a \vee f) \quad \text{and} \quad (c \wedge f) \leq c \wedge (a \vee f),$$

which by (ii) amounts to

$$b \leq c \quad \text{and} \quad b \leq a \vee f \quad \text{and} \quad c \wedge f \leq c \quad \text{and} \quad c \wedge f \leq a \vee f.$$

⁴For unordered sets P this has been proven by P. M. Whitman in 1941, although J. B. Nation states that Skolem achieved essentially the same in 1920. The natural extension to posets $(P, \leq)$ is due to R. P. Dilworth 1945. Reproving a 1964 result of R. A. Dean, H. Lakser [8] dealt with the lattice freely generated by $(P, \leq)$ and additionally preserving certain finite joins and meets in $(P, \leq)$.

Now by the converse of (iii), $b \not\leq a \vee f$ since $b \not\leq a, b \not\leq f$ in P . Therefore also $b \vee (c \wedge f) \not\leq c \wedge (a \vee f)$. For our $(P, \leq)$ of figure 3(a), $FL(P, \leq)$ happens to be finite and is depicted in figure 4. Observe that the "bubbles" in figure 4 are the congruence classes of the epimorphism $FL(P, \leq) \rightarrow FD(P, \leq)$ where $FD(P, \leq)$ is the free distributive lattice discussed in section 7. The lattices $FL(P, \leq)$ will recur in section 6 of Part II.




Figure 4:

7 The variety of distributive lattices

If $\mathcal{V}$ is the variety $\mathcal{D}$ of distributive lattices, then $F\mathcal{V}(P, \leq)$ will be written as $FD(P, \leq)$. Like every distributive lattice, $FD(P, \leq)$ is a subdirect product of two element factor lattices $D_2 = \{0, 1\}$. These factors can be neatly distinguished by the set of elements $\mathcal{F} \subseteq P$ that map upon 1 (as opposed to 0) under the projection $\pi : FD(P, \leq) \rightarrow D_2$. Since π is monotone, $\mathcal{F} = \pi^{-1}(1)$ is a (nonempty) order filter of $(P, \leq)$. All order filters $\mathcal{F}$ arise in this way by the universal mapping property of $FD(P, \leq)$. Thus we find that $FD(P, \leq)$ will have these 12 subdirectly irreducible factors:




Figure 5:

At this point we are stuck because the information provided by the " P -labellings" in Fig.5 is not enough to construct $FD(P, \leq)$. What one needs are the *connection maps* between any two P -labellings. We shall persue this line of thought in our sequel paper which deals with arbitrary finitely generated varieties of lattices (of which $\mathcal{D}$ is the simplest example). But here we tackle $FD(P, \leq)$ in a way that avoids subdirect products. The core is contained in the next lemma.

Lemma 1 *Let F and T be distributive lattices such that F is finite and $J_0(F) := J(F) \cup \{0\}$ is closed under meets. Then each $\wedge$ -preserving map $\bar{\phi} : J_0(F) \rightarrow T$ can be extended to a homomorphism $\Phi : F \rightarrow T$.*

Proof: For all $x \in F$ put

$$\Phi(x) := \bigvee \{\bar{\phi}(s) \mid s \leq x\}$$

where by convention s (or t or r) ranges over $J_0(F)$. Because $\bar{\phi}$ is monotone, Φ extends $\bar{\phi}$ and is monotone itself. It thus remains to show that Φ preserves $\wedge$ and $\vee$. As to $\wedge$, by assumption $s, t \in J_0(F)$ implies $s \wedge t \in J_0(F)$, and $\bar{\phi}(s) \wedge \bar{\phi}(t) = \bar{\phi}(s \wedge t)$. Hence

$$\begin{aligned} \Phi(x) \wedge \Phi(y) &= \bigvee \{\bar{\phi}(s) \mid s \leq x\} \wedge \bigvee \{\bar{\phi}(t) \mid t \leq y\} \\ &= \bigvee \{\bar{\phi}(s) \wedge \bar{\phi}(t) \mid s \leq x, t \leq y\} && \text{by distributivity} \\ &= \bigvee \{\bar{\phi}(s \wedge t) \mid s \leq x, t \leq y\} \\ &= \bigvee \{\bar{\phi}(r) \mid r \leq x \wedge y\} = \Phi(x \wedge y) \end{aligned}$$

By distributivity each $s \in J_0(F)$ is join-prime, that is, $s \leq x \vee y$ implies $s \leq x$ or $s \leq y$. Hence

$$\begin{aligned}
\Phi(x \vee y) &= \bigvee \{\bar{\phi}(s) \mid s \leq x \vee y\} \\
&= \bigvee \{\bar{\phi}(s) \mid s \leq x \text{ or } s \leq y\} \\
&\leq \bigvee \{\bar{\phi}(s) \mid s \leq x\} \vee \bigvee \{\bar{\phi}(s) \mid s \leq y\} \\
&= \Phi(x) \vee \Phi(y)
\end{aligned}$$

The inequality $\geq$ is trivial. ■

Theorem 3 *The lattice $FD(P, \leq)$ freely generated by the finite poset $(P, \leq)$ within the variety of all distributive lattices is isomorphic to $F_\vee(F_\wedge(P, \leq), \leq)$.*

Proof: For elements p, q of any poset, if the meet $p \wedge q$ happens to exist, then one checks that

$$(p \wedge q) \downarrow = p \downarrow \cap q \downarrow.$$

We view the $\wedge$ -semilattice $J_0 := F_\wedge(P, \leq)$ as a poset with smallest element 0. By Corollary 1 the $\cup$ -semilattice of nonempty order ideals of J_0 can be identified with $F := F_\vee(J_0, \leq)$. Since F has a smallest element (corresponding to $\{0\}$), it is a distributive lattice with $J_0(F)$ equal to J_0 . By the above remark J_0 is closed under meets.

Let now T be any distributive lattice and $\phi : P \rightarrow T$ a monotone map. Since J_0 is the free $\wedge$ -semilattice generated by $(P, \leq)$, ϕ can be extended to a $\wedge$ -preserving map $\bar{\phi}$ on J_0 . By Lemma 1, $\bar{\phi}$ further extends to a homomorphism $\Phi : F \rightarrow T$. ■

The proof given here is believed to be new. Theorem 3 is a special case of more general (but clumsier) results, e.g. by W. R. Tunnicliffe [9] or Yongming Li [10].

Example 5 Consider the poset $(P, \leq)$ from Fig.3(a). One checks that the fat subset J_0 of join irreducibles (including 0) of $FD(P, \leq)$ is a meet subsemilattice, and it is isomorphic to $F_\wedge(P, \leq)$ from Fig.3(c). The elements $a, \dots, g$ of P correspond to the doubly-irreducible elements of $FD(P, \leq)$:




Figure 6:

The subdirectly irreducible factors $\pi : D \rightarrow D_2$ of every finite distributive lattice D bijectively correspond to the join irreducibles q in that

$$\pi^{-1}(1) = \{x \in D \mid x \geq q\} \quad \text{and} \quad \pi^{-1}(0) = \{x \in D \mid x \not\geq q\}. \quad (15)$$

If for $D = FD(P, \leq)$ we take (say) $q = b$, we have $\pi^{-1}(1) \cap P = \{b, c, d, g\}$ and $\pi^{-1}(0) \cap P = \{a, e, f\}$ corresponds indeed to the eighth P -labelling in Fig.5.

Open problem: Set up an algorithm that computes *arbitrary* finitely presented lattices within $\mathcal{D}$, such as the free distributive lattice generated by a, b, c, d and subject to say $a \vee b = (c \wedge d) \vee (a \wedge b)$.

8 The variety of Boolean algebras

The class of all Boolean lattices is no variety since it is not closed under taking sublattices. But it becomes a variety if $0, 1$ are elevated to nullary operations (constants) and complementation $x \mapsto \bar{x}$ is added as unary operation. One then speaks of Boolean *algebras* and accordingly the definition in section 6 has to be adjusted to the extent that for the *free Boolean algebra* $FB(P, \leq)$ generated by the poset $(P, \leq)$, the homomorphism Φ in (b) must be a *Boolean* homomorphism in the sense that $\Phi(0) = 0$, $\Phi(1) = 1$, and $\Phi(\bar{x}) = \overline{\Phi(x)}$ for all $x \in FB(P, \leq)$. This section is based on [6, p. 107].

In order to describe $FB(P, \leq)$ for finite P , observe that any Boolean algebra $B(x_1, \dots, x_s)$ which is generated by s elements of some comprising Boolean algebra, by distributivity

and De Morgan's laws equals

$$B(x_1, \dots, x_s) = \left\{ \bigvee_{\delta \in I} (x_1^{\delta_1} \wedge x_2^{\delta_2} \wedge \dots \wedge x_s^{\delta_s}) \mid I \subseteq \{0, 1\}^s \right\}. \quad (16)$$

Here $\delta = (\delta_1, \delta_2, \dots, \delta_s)$ is a 0, 1-vector and $x_i^1 := x_i$, $x_i^0 := \overline{x_i}$. From (16) we conclude that the t atoms of $B(x_1, \dots, x_s)$ are exactly the *nonzero* elements among the at most 2^s elements $x_1^{\delta_1} \wedge \dots \wedge x_s^{\delta_s}$. It follows that if $FB(s)$ is the free Boolean algebra generated by s unordered elements, then

$$|FB(s)| \leq 2^{(2^s)}. \quad (17)$$

Example 6 Let $X := \{a, b, \dots, i\}$. What is the number t of atoms of the Boolean algebra $B(A_1, A_2, A_3, A_4) \subseteq \mathcal{P}(X)$ generated by $A_1 := \{a, c, d, f\}$, $A_2 := \{b, e, f, i\}$, $A_3 := \{a, b, c, d, e, g, i\}$ and $A_4 := \{e, g, h\}$?

	a	b	c	d	e	f	g	h	i
$A_1 =$	1	0	1	1	0	1	0	0	0
$A_2 =$	0	1	0	0	1	1	0	0	1
$A_3 =$	1	1	1	1	1	0	1	0	1
$A_4 =$	0	0	0	0	1	0	1	1	0

Here the characteristic vectors of the subsets $A_i \subseteq S$ are listed as the rows of a 4×9 matrix. Then the number t of equivalence classes of equal columns obviously is the number of $(\delta_1, \delta_2, \delta_3, \delta_4)$ with $A_1^{\delta_1} \cap A_2^{\delta_2} \cap A_3^{\delta_3} \cap A_4^{\delta_4} \neq \emptyset$. For instance, the three equal columns labelled by a, c, d yield $(\delta_1, \delta_2, \delta_3, \delta_4) = (1, 0, 1, 0)$ and $A_1^1 \cap A_2^0 \cap A_3^1 \cap A_4^0 = \{a, c, d\} \neq \emptyset$. In our case $t = 6$ whence $|B(A_1, A_2, A_3, A_4)| = 2^6 = 64$. $\square$

Picking s random sets from a r -set amounts to pick r random 0, 1-columns of length s . In Example 6 we had $r = 9$ and $s = 4$. As we saw, had the columns be distinct (which doesn't imply distinct rows), then $A_1, \dots, A_s$ would have generated $\mathcal{P}(X)$. The probability of getting distinct columns is easily calculated:

Theorem 4 *Let X be a r -set. Pick s (not necessarily distinct) sets $A_i \in \mathcal{P}(X)$ at random. The probability that $B(A_1, \dots, A_s)$ equals $\mathcal{P}(X)$ is*

$$\frac{2^s - 1}{2^s} \dots \frac{2^s - 2}{2^s} \dots \frac{2^s - (r - 1)}{2^s}$$

For instance, the probability that $s = 5$ random subsets $A_i \subseteq X := \{1, 2, 3, 4, 5\}$ generate the whole powerset is 0.72. When all $A_i \subseteq \{1, 2, \dots, 10\}$, the probability is 0.21. When $r = 2^s$, the probability for $|B(A_1, \dots, A_s)| = |\mathcal{P}(X)| = 2^{(2^s)}$ is still > 0 . In fact, in view of (17) such a $B(A_1, \dots, A_s)$ must be isomorphic to $FB(s)$. Let us now turn from s -element antichains to general posets $(P, \leq)$.

Theorem 5 *If $(P, \leq)$ is finite, then $FB(P, \leq)$ has exactly t atoms, where t is the number of order filters $\emptyset \subseteq F \subseteq P$.*

Proof: Assume w.l.o.g. $(P, \leq)$ is $(\{1, 2, \dots, s\}, \leq)$ and that $(\{A_1, \dots, A_s\}, \subseteq)$ is any fixed set system such that $i \leq j \Leftrightarrow A_i \subseteq A_j$ for all $i, j \in P$. In view of Example 6 observe that $A_1^{\delta_1} \cap \dots \cap A_s^{\delta_s}$ can be nonempty only if “ $\delta_i = 1 \Rightarrow \delta_j = 1$ ” whenever $i \leq j$. Hence the number of vectors $\delta = (\delta_1, \dots, \delta_s)$ with $A_1^{\delta_1} \cap \dots \cap A_s^{\delta_s} \neq \emptyset$ is *at most* the number t of order filters of P . On the other hand, choosing $(\{A_1, \dots, A_s\}, \subseteq)$ appropriately (as in Example 8 below), one can indeed obtain t nonempty sets $A_1^{\delta_1} \cap \dots \cap A_s^{\delta_s}$. This Boolean algebra of largest cardinality generated by $(P, \leq)$ can only be $FB(P, \leq)$ ■

Example 7 What is the size of the free Boolean algebra $FB(P, \leq)$ generated by our companion poset $(P, \leq)$ of figure 3(a)? As opposed to Example 6, here the 0, 1-matrix is generated by concatenating the *columns*. Namely, if we list the characteristic vectors of the $t = 14$ order filters of $(P, \leq)$ as the (distinct) columns of a 7×14 matrix, then the sets corresponding to the rows yield a set system isomorphic to $(P, \leq)$ (check), and all intersections $A_a^{\delta_a} \cap \dots \cap A_g^{\delta_g}$, where δ is a column of the matrix, are nonempty (as argued in Example 7). Thus $|FB(P, \leq)| = 2^{14} = 16384$.

	$\emptyset$	$a \uparrow$	$b \uparrow$	$c \uparrow$	$d \uparrow$	$e \uparrow$	$f \uparrow$	$g \uparrow$	$a, e \uparrow$	$a, f \uparrow$	$b, f \uparrow$	$c, f \uparrow$	$c, g \uparrow$	$d, g \uparrow$
$A_a =$	0	1	0	0	0	0	0	0	1	1	0	0	0	0
$A_b =$	0	1	1	0	0	1	0	0	1	1	1	0	0	0
$A_c =$	0	1	1	1	0	1	0	0	1	1	1	1	1	0
$A_d =$	0	1	1	1	1	1	1	0	1	1	1	1	1	1
$A_e =$	0	0	0	0	0	1	0	0	1	0	0	0	0	0
$A_f =$	0	0	0	0	0	1	1	0	1	1	1	1	0	0
$A_g =$	0	1	1	0	0	1	1	1	1	1	1	1	1	1

References

- [1] V. Duquenne, *The core of finite lattices*, Discrete Mathematics 88 (1991) 133-147.
- [2] G. Grätzer, *General Lattice Theory*, Birkhäuser Verlag, Basel 2003.
- [3] J.Y. Semegni, *On the computation of freely generated modular lattices*, Ph.D. Thesis, University of Stellenbosch, 2008.
- [4] M. Wild, *Halbgruppen und formale Sprachen*, lecture notes, 163 pages, 1997.
- [5] M. Wild, *Computing various types of lattices freely generated by posets*, Note di matematica e fisica 10 (1999) 99-128.
- [6] M. Wild, *Generating all satisfying truth assignments for Horn-SAT*, in preparation.
- [7] R. Wille, *On lattices freely generated by finite partially ordered sets*, Coll. Math. Soc. Janos Bolyai 1975, p.581-593.

- [8] H. Lakser, *Lattices freely generated by an order and preserving certain bounds*, submitted.
- [9] W.R. Tunnicliffe, *The free completely distributive lattice over a poset*, Algebra Universalis 21 (1985) 133-135.
- [10] Yongming Li, *Generalized (S, I) -complete free completely distributive lattices generated by posets*, Semigroup Forum 57 (1998) 240-248.